

# Ordinary Thinking

## Objective Questions

### Pressure and Density

1. If pressure at half the depth of a lake is equal to  $\frac{2}{3}$  pressure at the bottom of the lake then what is the depth of the lake

[RPET 2000]

- (a) 10 m                      (b) 20 m  
(c) 60 m                      (d) 30 m

2. Two bodies are in equilibrium when suspended in water from the arms of a balance. The mass of one body is 36 g and its density is  $9 \text{ g/cm}^3$ . If the mass of the other is 48 g, its density in  $\text{g/cm}^3$  is

[CBSE PMT 1994]


- (a)  $\frac{4}{3}$                           (b)  $\frac{3}{2}$   
(c) 3                              (d) 5

3. An inverted bell lying at the bottom of a lake 47.6 m deep has  $50 \text{ cm}^3$  of air trapped in it. The bell is brought to the surface of the lake. The volume of the trapped air will be (atmospheric pressure = 70 cm of Hg and density of Hg =  $13.6 \text{ g/cm}^3$ )

[CPMT 1989]


- (a)  $350 \text{ cm}^3$                       (b)  $300 \text{ cm}^3$   
(c)  $250 \text{ cm}^3$                       (d)  $22 \text{ cm}^3$

4. A uniformly tapering vessel is filled with a liquid of density  $900 \text{ kg/m}^3$ . The force that acts on the base of the vessel due to the liquid is ( $g = 10 \text{ ms}^{-2}$ )



- (a) 3.6 N  
(b) 7.2 N  
(c) 9.0 N  
(d) 14.4 N

5. A siphon in use is demonstrated in the following figure. The density of the liquid flowing in siphon is  $1.5 \text{ gm/cc}$ . The pressure difference between the point P and S will be



- (a)  $10^5 \text{ N/m}$   
(b)  $2 \times 10^5 \text{ N/m}$   
(c) Zero  
(d) Infinity

6. The height of a mercury barometer is 75 cm at sea level and 50 cm at the top of a hill. Ratio of density of mercury to that of air is  $10^4$ . The height of the hill is

- (a) 250 m                      (b) 2.5 km  
(c) 1.25 km                      (d) 750 m

7. Density of ice is  $\rho$  and that of water is  $\sigma$ . What will be the decrease in volume when a mass  $M$  of ice melts

- (a)  $\frac{M}{\sigma - \rho}$                       (b)  $\frac{\sigma - \rho}{M}$   
(c)  $M \left[ \frac{1}{\rho} - \frac{1}{\sigma} \right]$                       (d)  $\frac{1}{M} \left[ \frac{1}{\rho} - \frac{1}{\sigma} \right]$

8. Equal masses of water and a liquid of density 2 are mixed together, then the mixture has a density of

- (a)  $\frac{2}{3}$                               (b)  $\frac{4}{3}$   
(c)  $\frac{3}{2}$                               (d) 3

9. A body of density  $d_1$  is counterpoised by  $Mg$  of weights of density  $d_2$  in air of density  $d$ . Then the true mass of the body is

- (a)  $M$                               (b)  $M \left( 1 - \frac{d}{d_2} \right)$   
(c)  $M \left( 1 - \frac{d}{d_1} \right)$                       (d)  $\frac{M(1 - d/d_2)}{(1 - d/d_1)}$

10. The pressure at the bottom of a tank containing a liquid does not depend on

[Kerala (Engg.) 2002]


- (a) Acceleration due to gravity  
(b) Height of the liquid column  
(c) Area of the bottom surface  
(d) Nature of the liquid

11. When a large bubble rises from the bottom of a lake to the surface. Its radius doubles. If atmospheric pressure is equal to that of column of water height  $H$ , then the depth of lake is

[AIIMS 1995; AFMC 1997]

- (a)  $H$                               (b)  $2H$


- (c)  $7H$  (d)  $8H$
12. The volume of an air bubble becomes three times as it rises from the bottom of a lake to its surface. Assuming atmospheric pressure to be  $75 \text{ cm}$  of  $Hg$  and the density of water to be  $1/10$  of the density of mercury, the depth of the lake is [AMU 1995]
- (a)  $5 \text{ m}$  (b)  $10 \text{ m}$   
(c)  $15 \text{ m}$  (d)  $20 \text{ m}$
13. The value of  $g$  at a place decreases by  $2\%$ . The barometric height of mercury
- (a) Increases by  $2\%$   
(b) Decreases by  $2\%$   
(c) Remains unchanged  
(d) Sometimes increases and sometimes decreases
14. A barometer kept in a stationary elevator reads  $76 \text{ cm}$ . If the elevator starts accelerating up the reading will be
- (a) Zero (b) Equal to  $76 \text{ cm}$   
(c) More than  $76 \text{ cm}$  (d) Less than  $76 \text{ cm}$
15. A closed rectangular tank is completely filled with water and is accelerated horizontally with an acceleration  $a$  towards right. Pressure is (i) maximum at, and (ii) minimum at



- (a) (i) B (ii) D  
(b) (i) C (ii) D  
(c) (i) B (ii) C  
(d) (i) B (ii) A
16. A beaker containing a liquid is kept inside a big closed jar. If the air inside the jar is continuously pumped out, the pressure in the liquid near the bottom of the liquid will
- (a) Increases  
(b) Decreases  
(c) Remain constant  
(d) First decrease and then increase
17. A barometer tube reads  $76 \text{ cm}$  of mercury. If the tube is gradually inclined at an angle of  $60^\circ$  with vertical, keeping the open end immersed

in the mercury reservoir, the length of the mercury column will be

- (a)  $152 \text{ cm}$  (b)  $76 \text{ cm}$   
(c)  $38 \text{ cm}$  (d)  $38\sqrt{3} \text{ cm}$
18. The height to which a cylindrical vessel be filled with a homogeneous liquid, to make the average force with which the liquid presses the side of the vessel equal to the force exerted by the liquid on the bottom of the vessel, is equal to
- (a) Half of the radius of the vessel  
(b) Radius of the vessel  
(c) One-fourth of the radius of the vessel  
(d) Three-fourth of the radius of the vessel
19. A vertical U-tube of uniform inner cross section contains mercury in both sides of its arms. A glycerin (density =  $1.3 \text{ g/cm}^3$ ) column of length  $10 \text{ cm}$  is introduced into one of its arms. Oil of density  $0.8 \text{ gm/cm}^3$  is poured into the other arm until the upper surfaces of the oil and glycerin are in the same horizontal level. Find the length of the oil column, Density of mercury =  $13.6 \text{ g/cm}^3$



- (a)  $10.4 \text{ cm}$   
(b)  $8.2 \text{ cm}$   
(c)  $7.2 \text{ cm}$   
(d)  $9.6 \text{ cm}$
20. A triangular lamina of area  $A$  and height  $h$  is immersed in a liquid of density  $\rho$  in a vertical plane with its base on the surface of the liquid. The thrust on the lamina is
- (a)  $\frac{1}{2} A\rho gh$  (b)  $\frac{1}{3} A\rho gh$   
(c)  $\frac{1}{6} A\rho gh$  (d)  $\frac{2}{3} A\rho gh$
21. If two liquids of same masses but densities  $\rho_1$  and  $\rho_2$  respectively are mixed, then density of mixture is given by
- (a)  $\rho = \frac{\rho_1 + \rho_2}{2}$  (b)  $\rho = \frac{\rho_1 + \rho_2}{2\rho_1\rho_2}$

(c)  $\rho = \frac{2\rho_1\rho_2}{\rho_1 + \rho_2}$       (d)  $\rho = \frac{\rho_1\rho_2}{\rho_1 + \rho_2}$

22. If two liquids of same volume but different densities  $\rho_1$  and  $\rho_2$  are mixed, then density of mixture is given by

(a)  $\rho = \frac{\rho_1 + \rho_2}{2}$       (b)  $\rho = \frac{\rho_1\rho_2}{2\rho_1\rho_2}$

(c)  $\rho = \frac{2\rho_1\rho_2}{\rho_1 + \rho_2}$       (d)  $\rho = \frac{\rho_1\rho_2}{\rho_1 + \rho_2}$

23. The density  $\rho$  of water of bulk modulus  $B$  at a depth  $y$  in the ocean is related to the density at surface  $\rho_0$  by the relation

(a)  $\rho = \rho_0 \left[ 1 - \frac{\rho_0 g y}{B} \right]$       (b)  $\rho = \rho_0 \left[ 1 + \frac{\rho_0 g y}{B} \right]$

(c)  $\rho = \rho_0 \left[ 1 + \frac{B}{\rho_0 h g y} \right]$       (d)  $\rho = \rho_0 \left[ 1 - \frac{B}{\rho_0 g y} \right]$

24. With rise in temperature, density of a given body changes according to one of the following relations

(a)  $\rho = \rho_0 [1 + \gamma d\theta]$       (b)  $\rho = \rho_0 [1 - \gamma d\theta]$

(c)  $\rho = \rho_0 \gamma d\theta$       (d)  $\rho = \rho_0 / \gamma d\theta$

25. Three liquids of densities  $d, 2d$  and  $3d$  are mixed in equal volumes. Then the density of the mixture is

(a)  $d$       (b)  $2d$


(c)  $3d$       (d)  $5d$

26. Three liquids of densities  $d, 2d$  and  $3d$  are mixed in equal proportions of weights. The relative density of the mixture is

(a)  $\frac{11d}{7}$       (b)  $\frac{18d}{11}$

(c)  $\frac{13d}{9}$       (d)  $\frac{23d}{18}$

27. From the adjacent figure, the correct observation is




(a) The pressure on the bottom of tank (a) is greater than at the bottom of (b).

(b) The pressure on the bottom of the tank (a) is smaller than at the bottom of (b)

(c) The pressure depend on the shape of the container

(d) The pressure on the bottom of (a) and (b) is the same

28. A given shaped glass tube having uniform cross section is filled with water and is mounted on a rotatable shaft as shown in figure. If the tube is rotated with a constant angular velocity  $\omega$  then



(a) Water levels in both sections A and B go up

(b) Water level in Section A goes up and that in B comes down

(c) Water level in Section A comes down and that in B it goes up

(d) Water levels remains same in both sections

29. Why the dam of water reservoir is thick at the bottom

[AFMC 2005]

(a) Quantity of water increases with depth

(b) Density of water increases with depth

(c) Pressure of water increases with depth

(d) Temperature of water increases with depth

30. Air is blown through a hole on a closed pipe containing liquid. Then the pressure will

[AFMC 2005]

(a) Increase on sides

(b) Increase downwards

(c) Increase in all directions

(d) Never increases

31. Radius of an air bubble at the bottom of the lake is  $r$  and it becomes  $2r$  when the air bubbles rises to the top surface of the lake. If  $P$  cm of water be the atmospheric pressure, then the depth of the lake is

[Kerla PET 2005]

- (a)  $2p$  (b)  $8p$   
(c)  $4p$  (d)  $7p$

Pascal's Law and Archimides Principle

1. An ice berg of density  $900 \text{ Kg/m}^3$  is floating in water of density  $1000 \text{ Kg/m}^3$ . The percentage of volume of ice-cube outside the water is

[CPMT 2004]

- (a) 20% (b) 35%  
(c) 10% (d) 25%

2. A log of wood of mass  $120 \text{ Kg}$  floats in water. The weight that can be put on the raft to make it just sink, should be (density of wood =  $600 \text{ Kg/m}^3$ )

[CPMT 2004]

- (a)  $80 \text{ Kg}$  (b)  $50 \text{ Kg}$   
(c)  $60 \text{ Kg}$  (d)  $30 \text{ Kg}$

3. A hemispherical bowl just floats without sinking in a liquid of density  $1.2 \times 10^3 \text{ kg/m}^3$ . If outer diameter and the density of the bowl are  $1 \text{ m}$  and  $2 \times 10^4 \text{ kg/m}^3$  respectively, then the inner diameter of the bowl will be

[SCRA 1998]

- (a)  $0.94 \text{ m}$  (b)  $0.97 \text{ m}$   
(c)  $0.98 \text{ m}$  (d)  $0.99 \text{ m}$

4. In making an alloy, a substance of specific gravity  $s_1$  and mass  $m_1$  is mixed with another substance of specific gravity  $s_2$  and mass  $m_2$ ; then the specific gravity of the alloy is

[CPMT 1995]

- (a)  $\left(\frac{m_1 + m_2}{s_1 + s_2}\right)$  (b)  $\left(\frac{s_1 s_2}{m_1 + m_2}\right)$

- (c)  $\frac{m_1 + m_2}{\left(\frac{m_1}{s_1} + \frac{m_2}{s_2}\right)}$  (d)  $\frac{\left(\frac{m_1}{s_1} + \frac{m_2}{s_2}\right)}{m_1 + m_2}$

5. A concrete sphere of radius  $R$  has a cavity of radius  $r$  which is packed with sawdust. The specific gravities of concrete and sawdust are respectively  $2.4$  and  $0.3$  for this sphere to float with its entire volume submerged under water. Ratio of mass of concrete to mass of sawdust will be

[AIIMS 1995]

- (a)  $8$  (b)  $4$   
(c)  $3$  (d) Zero

6. A metallic block of density  $5 \text{ gm cm}^{-3}$  and having dimensions  $5 \text{ cm} \times 5 \text{ cm} \times 5 \text{ cm}$  is weighed in water. Its apparent weight will be

- (a)  $5 \times 5 \times 5 \times 5 \text{ gf}$  (b)  $4 \times 4 \times 4 \times 4 \text{ gf}$   
(c)  $5 \times 4 \times 4 \times 4 \text{ gf}$  (d)  $4 \times 5 \times 5 \times 5 \text{ gf}$


7. A cubical block is floating in a liquid with half of its volume immersed in the liquid. When the whole system accelerates upwards with acceleration of  $g/3$ , the fraction of volume immersed in the liquid will be

(a)  $\frac{1}{2}$

(b)  $\frac{3}{8}$

(c)  $\frac{2}{3}$

(d)  $\frac{3}{4}$



8. A silver ingot weighing  $2.1 \text{ kg}$  is held by a string so as to be completely immersed in a liquid of relative density  $0.8$ . The relative density of silver is  $10.5$ . The tension in the string in  $\text{kg-wt}$  is

- (a)  $1.6$  (b)  $1.94$   
(c)  $3.1$  (d)  $5.25$


9. A sample of metal weighs  $210 \text{ gm}$  in air,  $180 \text{ gm}$  in water and  $120 \text{ gm}$  in liquid. Then relative density (RD) of

- (a) Metal is  $3$  (b) Metal is  $7$   
(c) Liquid is  $3$  (d) Liquid is  $\frac{1}{3}$

10. Two solids  $A$  and  $B$  float in water. It is observed that  $A$  floats with half its volume immersed and  $B$  floats with  $2/3$  of its volume immersed. Compare the densities of  $A$  and  $B$

- (a)  $4 : 3$  (b)  $2 : 3$   
(c)  $3 : 4$  (d)  $1 : 3$


11. The fraction of a floating object of volume  $V_0$  and density  $d_0$  above the surface of a liquid of density  $d$  will be

- (a)  $\frac{d_0}{d}$  (b)  $\frac{dd_0}{d+d_0}$   
 (c)  $\frac{d-d_0}{d}$  (d)  $\frac{dd_0}{d-d_0}$
12. Pressure applied to an enclosed fluid is transmitted undiminished to every portion of the fluid and the walls of the containing vessel. This law was first formulated by  
 (a) Bernoulli (b) Archimedes  
 (c) Boyle (d) Pascal
13. A block of steel of size  $5\text{ cm} \times 5\text{ cm} \times 5\text{ cm}$  is weighed in water. If the relative density of steel is 7, its apparent weight is  
 [AFMC 1997]  
 (a)  $6 \times 5 \times 5 \times 5\text{ gf}$  (b)  $4 \times 4 \times 4 \times 7\text{ gf}$   
 (c)  $5 \times 5 \times 5 \times 7\text{ gf}$  (d)  $4 \times 4 \times 4 \times 6\text{ gf}$
14. A body is just floating on the surface of a liquid. The density of the body is same as that of the liquid. The body is slightly pushed down. What will happen to the body [AIIMS 1980]  
 (a) It will slowly come back to its earlier position  
 (b) It will remain submerged, where it is left  
 (c) It will sink  
 (d) It will come out violently
15. A cork is submerged in water by a spring attached to the bottom of a bowl. When the bowl is kept in an elevator moving with acceleration downwards, the length of spring  
 (a) Increases (b) Decreases  
 (c) Remains unchanged (d) None of these
16. A solid sphere of density  $\eta (> 1)$  times lighter than water is suspended in a water tank by a string tied to its base as shown in fig. If the mass of the sphere is  $m$  then the tension in the string is given by  
 (a)  $\left(\frac{\eta-1}{\eta}\right)mg$  (b)  $\eta mg$   
 (c)  $\frac{mg}{\eta-1}$  (d)  $(\eta-1)mg$
- 
17. A hollow sphere of volume  $V$  is floating on water surface with *half* immersed in it. What should be the minimum volume of water poured inside the sphere so that the sphere now sinks into the water  
 (a)  $V/2$  (b)  $V/3$   
 (c)  $V/4$  (d)  $V$
18. A rectangular block is  $5\text{ cm} \times 5\text{ cm} \times 10\text{ cm}$  in size. The block is floating in water with  $5\text{ cm}$  side vertical. If it floats with  $10\text{ cm}$  side vertical, what change will occur in the level of water?  
 (a) No change  
 (b) It will rise  
 (c) It will fall  
 (d) It may rise or fall depending on the density of block
19. A ball whose density is  $0.4 \times 10^3\text{ kg/m}^3$  falls into water from a height of  $9\text{ cm}$ . To what depth does the ball sink  
 (a)  $9\text{ cm}$  (b)  $6\text{ cm}$   
 (c)  $4.5\text{ cm}$  (d)  $2.25\text{ cm}$
20. Two solids  $A$  and  $B$  float in water. It is observed that  $A$  floats with  $\frac{1}{2}$  of its body immersed in water and  $B$  floats with  $\frac{1}{4}$  of its volume above the water level. The ratio of the density of  $A$  to that of  $B$  is  
 (a)  $4 : 3$  (b)  $2 : 3$   
 (c)  $3 : 4$  (d)  $1 : 2$
21. A boat carrying steel balls is floating on the surface of water in a tank. If the balls are thrown into the tank one by one, how will it affect the level of water [J&K CET 2005]  
 (a) It will remain unchanged  
 (b) It will rise  
 (c) It will fall  
 (d) First it will rise and then fall
22. Two pieces of metal when immersed in a liquid have equal upthrust on them; then  
 (a) Both pieces must have equal weights

Fluid Flow


- (b) Both pieces must have equal densities  
 (c) Both pieces must have equal volumes  
 (d) Both are floating to the same depth
23. A wooden cylinder floats vertically in water with half of its length immersed. The density of wood is
- (a) Equal of that of water  
 (b) Half the density of water  
 (c) Double the density of water  
 (d) The question is incomplete

24. A candle of diameter  $d$  is floating on a liquid in a cylindrical container of diameter  $D$  ( $D \gg d$ ) as shown in figure. If it is burning at the rate of  $2\text{ cm/hour}$  then the top of the candle will




- (a) Remain at the same height  
 (b) Fall at the rate of  $1\text{ cm/hour}$   
 (c) Fall at the rate of  $2\text{ cm/hour}$   
 (d) Go up the rate of  $1\text{ cm/hour}$
25. An ice block contains a glass ball when the ice melts within the water containing vessel, the level of water [AFMC 2005]
- (a) Rises (b) Falls  
 (c) Unchanged (d) First rises and then falls
26. A large ship can float but a steel needle sinks because of [AFMC 2005]
- (a) Viscosity (b) Surface tension  
 (c) Density (d) None of these
27. Construction of submarines is based on [Kerala PMT 2005]
- (a) Archimedes' principle (b) Bernoulli's theorem  
 (c) Pascal's law (d) Newton's laws

1. In which one of the following cases will the liquid flow in a pipe be most streamlined [Pb. CET 2005]
- (a) Liquid of high viscosity and high density flowing through a pipe of small radius  
 (b) Liquid of high viscosity and low density flowing through a pipe of small radius  
 (c) Liquid of low viscosity and low density flowing through a pipe of large radius  
 (d) Liquid of low viscosity and high density flowing through a pipe of large radius
2. Two water pipes of diameters  $2\text{ cm}$  and  $4\text{ cm}$  are connected with the main supply line. The velocity of flow of water in the pipe of  $2\text{ cm}$  diameter is [MNR 1980]
- (a) 4 times that in the other pipe  
 (b)  $\frac{1}{4}$  times that in the other pipe  
 (c) 2 times that in the other pipe  
 (d)  $\frac{1}{2}$  times that in the other pipe
3. An incompressible liquid flows through a horizontal tube as shown in the following fig. Then the velocity  $v$  of the fluid is



- (a)  $3.0\text{ m/s}$  (b)  $1.5\text{ m/s}$   
 (c)  $1.0\text{ m/s}$  (d)  $2.25\text{ m/s}$
4. Water enters through end A with speed  $v_1$  and leaves through end B with speed  $v_2$  of a cylindrical tube AB. The tube is always completely filled with water. In case I tube is horizontal and in case II it is vertical with end A upwards and in case III it is vertical with end B upwards. We have  $v_1 = v_2$  for

- (a) Case I (b) Case II  
(c) Case III (d) Each case
5. Water is moving with a speed of  $5.18 \text{ ms}^{-1}$  through a pipe with a cross-sectional area of  $4.20 \text{ cm}^2$ . The water gradually descends  $9.66 \text{ m}$  as the pipe increase in area to  $7.60 \text{ cm}^2$ . The speed of flow at the lower level is  
(a)  $3.0 \text{ ms}^{-1}$  (b)  $5.7 \text{ ms}^{-1}$   
(c)  $3.82 \text{ ms}^{-1}$  (d)  $2.86 \text{ ms}^{-1}$
6. The velocity of kerosene oil in a horizontal pipe is  $5 \text{ m/s}$ . If  $g = 10 \text{ m/s}^2$  then the velocity head of oil will be  
(a)  $1.25 \text{ m}$  (b)  $12.5 \text{ m}$   
(c)  $0.125 \text{ m}$  (d)  $125 \text{ m}$
7. In the following fig. is shown the flow of liquid through a horizontal pipe. Three tubes  $A$ ,  $B$  and  $C$  are connected to the pipe. The radii of the tubes  $A$ ,  $B$  and  $C$  at the junction are respectively  $2 \text{ cm}$ ,  $1 \text{ cm}$  and  $2 \text{ cm}$ . It can be said that the



- (a) Height of the liquid in the tube  $A$  is maximum  
(b) Height of the liquid in the tubes  $A$  and  $B$  is the same  
(c) Height of the liquid in all the three tubes is the same  
(d) Height of the liquid in the tubes  $A$  and  $C$  is the same
8. A manometer connected to a closed tap reads  $3.5 \times 10^5 \text{ N/m}^2$ . When the valve is opened, the reading of manometer falls to  $3.0 \times 10^5 \text{ N/m}^2$ , then velocity of flow of water is  
(a)  $100 \text{ m/s}$  (b)  $10 \text{ m/s}$   
(c)  $1 \text{ m/s}$  (d)  $10\sqrt{10} \text{ m/s}$
9. Air is streaming past a horizontal air plane wing such that its speed in  $120 \text{ m/s}$  over the upper surface and  $90 \text{ m/s}$  at the lower surface.

If the density of air is  $1.3 \text{ kg per metre}^3$  and the wing is  $10 \text{ m}$  long and has an average width of  $2 \text{ m}$ , then the difference of the pressure on the two sides of the wing of

- (a)  $4095.0 \text{ Pascal}$  (b)  $409.50 \text{ Pascal}$   
(c)  $40.950 \text{ Pascal}$  (d)  $4.0950 \text{ Pascal}$
10. A large tank filled with water to a height ' $h$ ' is to be emptied through a small hole at the bottom. The ratio of time taken for the level of water to fall from  $h$  to  $\frac{h}{2}$  and from  $\frac{h}{2}$  to zero is  
[EAMCET (Engg.) 2003]  
(a)  $\sqrt{2}$  (b)  $\frac{1}{\sqrt{2}}$   
(c)  $\sqrt{2} - 1$  (d)  $\frac{1}{\sqrt{2} - 1}$
11. A cylinder of height  $20 \text{ m}$  is completely filled with water. The velocity of efflux of water (in  $\text{m/s}$ ) through a small hole on the side wall of the cylinder near its bottom is  
[AIEEE 2002]  
(a)  $10$  (b)  $20$   
(c)  $25.5$  (d)  $5$
12. There is a hole in the bottom of tank having water. If total pressure at bottom is  $3 \text{ atm}$  ( $1 \text{ atm} = 10^5 \text{ N/m}^2$ ) then the velocity of water flowing from hole is [CPMT 2002]  
(a)  $\sqrt{400} \text{ m/s}$  (b)  $\sqrt{600} \text{ m/s}$   
(c)  $\sqrt{60} \text{ m/s}$  (d) None of these
13. There is a hole of area  $A$  at the bottom of cylindrical vessel. Water is filled up to a height  $h$  and water flows out in  $t$  second. If water is filled to a height  $4h$ , it will flow out in time equal to [MP PMT 1997]  
(a)  $t$  (b)  $4t$   
(c)  $2t$  (d)  $t/4$
14. A cylindrical tank has a hole of  $1 \text{ cm}^2$  in its bottom. If the water is allowed to flow into the tank from a tube above it at the rate of  $70 \text{ cm}^3/\text{sec}$ . then the maximum height up to which water can rise in the tank is  
(a)  $2.5 \text{ cm}$  (b)  $5 \text{ cm}$

- (c) 10 cm (d) 0.25 cm
15. A square plate of 0.1 m side moves parallel to a second plate with a velocity of 0.1 m/s, both plates being immersed in water. If the viscous force is 0.002 N and the coefficient of viscosity is 0.01 poise, distance between the plates in m is

[EAMCET (Med.) 2003]

- (a) 0.1 (b) 0.05  
(c) 0.005 (d) 0.0005
16. Spherical balls of radius 'r' are falling in a viscous fluid of viscosity ' $\eta$ ' with a velocity 'v'. The retarding viscous force acting on the spherical ball is

[AIEEE 2004]

- (a) Inversely proportional to 'r' but directly proportional to velocity 'v'  
(b) Directly proportional to both radius 'r' and velocity 'v'  
(c) Inversely proportional to both radius 'r' and velocity 'v'  
(d) Directly proportional to 'r' but inversely proportional to 'v'

17. A small sphere of mass  $m$  is dropped from a great height. After it has fallen 100 m, it has attained its terminal velocity and continues to fall at that speed. The work done by air friction against the sphere during the first 100 m of fall is

[MP PMT 1990]


- (a) Greater than the work done by air friction in the second 100 m  
(b) Less than the work done by air friction in the second 100 m  
(c) Equal to 100 mg  
(d) Greater than 100 mg
18. Two drops of the same radius are falling through air with a steady velocity of 5 cm per sec. If the two drops coalesce, the terminal velocity would be

[MP PMT 1990]

- (a) 10 cm per sec (b) 2.5 cm per sec  
(c)  $5 \times (4)^{1/3}$  cm per sec (d)  $5 \times \sqrt{2}$  cm per sec

19. A ball of radius  $r$  and density  $\rho$  falls freely under gravity through a distance  $h$  before entering water. Velocity of ball does not change even on entering water. If viscosity of water is  $\eta$ , the value of  $h$  is given by

- (a)  $\frac{2}{9} r^2 \left( \frac{1-\rho}{\eta} \right) g$   
(b)  $\frac{2}{81} r^2 \left( \frac{\rho-1}{\eta} \right) g$   
(c)  $\frac{2}{81} r^4 \left( \frac{\rho-1}{\eta} \right)^2 g$   
(d)  $\frac{2}{9} r^4 \left( \frac{\rho-1}{\eta} \right)^2 g$



20. The rate of steady volume flow of water through a capillary tube of length ' $l$ ' and radius ' $r$ ' under a pressure difference of  $P$  is  $V$ . This tube is connected with another tube of the same length but half the radius in series. Then the rate of steady volume flow through them is (The pressure difference across the combination is  $P$ )

[EAMCET (Engg.) 2003]

- (a)  $\frac{V}{16}$  (b)  $\frac{V}{17}$   
(c)  $\frac{16V}{17}$  (d)  $\frac{17V}{16}$

21. A liquid is flowing in a horizontal uniform capillary tube under a constant pressure difference  $P$ . The value of pressure for which the rate of flow of the liquid is doubled when the radius and length both are doubled is

[EAMCET 2001]

- (a)  $P$  (b)  $\frac{3P}{4}$   
(c)  $\frac{P}{2}$  (d)  $\frac{P}{4}$

22. We have two (narrow) capillary tubes  $T_1$  and  $T_2$ . Their lengths are  $l_1$  and  $l_2$  and radii of cross-section are  $r_1$  and  $r_2$  respectively. The rate of flow of water under a pressure difference  $P$  through tube  $T_1$  is  $8 \text{ cm}^3/\text{sec}$ . If  $l_1 = 2l_2$  and  $r_1 = r_2$ , what will be the rate of flow when the two tubes are connected in series and pressure difference across the combination is same as before ( $= P$ )



- (a)  $4 \text{ cm}^3/\text{sec}$                       (b)  $(16/3) \text{ cm}^3/\text{sec}$   
 (c)  $(8/17) \text{ cm}^3/\text{sec}$                 (d) None of these
23. In a laminar flow the velocity of the liquid in contact with the walls of the tube is  
 (a) Zero  
 (b) Maximum  
 (c) In between zero and maximum  
 (d) Equal to critical velocity
24. In a turbulent flow, the velocity of the liquid molecules in contact with the walls of the tube is  
 (a) Zero  
 (b) Maximum  
 (c) Equal to critical velocity  
 (d) May have any value
25. The Reynolds number of a flow is the ratio of  
 (a) Gravity to viscous force  
 (b) Gravity force to pressure force  
 (c) Inertia forces to viscous force  
 (d) Viscous forces to pressure forces
26. Water is flowing through a tube of non-uniform cross-section ratio of the radius at entry and exit end of the pipe is 3 : 2. Then the ratio of velocities at entry and exit of liquid is  
 [RPMT 2001]  
 (a) 4 : 9                                      (b) 9 : 4  
 (c) 8 : 27                                    (d) 1 : 1
27. Water is flowing through a horizontal pipe of non-uniform cross-section. At the extreme narrow portion of the pipe, the water will have  
 [MP PMT 1992]  
 (a) Maximum speed and least pressure  
 (b) Maximum pressure and least speed  
 (c) Both pressure and speed maximum  
 (d) Both pressure and speed least
28. A liquid flows in a tube from left to right as shown in figure.  $A_1$  and  $A_2$  are the cross-sections of the portions of the tube as shown. Then the ratio of speeds  $v_1 / v_2$  will be



- (b)  $A_2 / A_1$   
 (c)  $\sqrt{A_2} / \sqrt{A_1}$   
 (d)  $\sqrt{A_1} / \sqrt{A_2}$
29. In a streamline flow  
 (a) The speed of a particle always remains same  
 (b) The velocity of a particle always remains same  
 (c) The kinetic energies of all the particles arriving at a given point are the same  
 (d) The moments of all the particles arriving at a given point are the same
30. An application of Bernoulli's equation for fluid flow is found in  
 [IIT-JEE (Screening) 1994]  
 (a) Dynamic lift of an aeroplane  
 (b) Viscosity meter  
 (c) Capillary rise  
 (d) Hydraulic press
31. The Working of an atomizer depends upon  
 [MP PMT 1992; AFMC 2005]  
 (a) Bernoulli's theorem (b) Boyle's law  
 (c) Archimedes principle (d)
32. The pans of a physical balance are in equilibrium. Air is blown under the right hand pan; then the right hand pan will  
 (a) Move up  
 (b) Move down  
 (c) Move erratically  
 (d) Remain at the same level
33. According to Bernoulli's equation

$$\frac{P}{\rho g} + h + \frac{1}{2} \frac{v^2}{g} = \text{constant}$$

The terms  $A$ ,  $B$  and  $C$  are generally called respectively:

- (a) Gravitational head, pressure head and velocity head

- (b) Gravity, gravitational head and velocity head
- (c) Pressure head, gravitational head and velocity head
- (d) Gravity, pressure and velocity head

34. At what speed the velocity head of a stream of water be equal to 40 cm of Hg


- (a) 282.8 cm/sec                      (b) 432.6 cm/sec
- (c) 632.6 cm/sec                    (d) 832.6 cm/sec

35. The weight of an aeroplane flying in air is balanced by

- (a) Upthrust of the air which will be equal to the weight of the air having the same volume as the plane
- (b) Force due to the pressure difference between the upper and lower surfaces of the wings, created by different air speeds on the surface
- (c) Vertical component of the thrust created by air currents striking the lower surface of the wings
- (d) Force due to the reaction of gases ejected by the revolving propeller

36. In this figure, an ideal liquid flows through the tube, which is of uniform cross-section. The liquid has velocities  $v_A$  and  $v_B$ , and pressure  $P_A$  and  $P_B$  at points A and B respectively


- (a)  $v_A = v_B$
- (b)  $v_B > v_A$
- (c)  $P_A = P_B$
- (d)  $P_B > P_A$



37. A liquid flows through a horizontal tube. The velocities of the liquid in the two sections, which have areas of cross-section  $A_1$  and  $A_2$ , are  $v_1$  and  $v_2$  respectively. The difference in the levels of the liquid in the two vertical tubes is  $h$

(a) The volume of the liquid flowing through the tube in unit time is

- (b)  $v_2 - v_1 = \sqrt{2gh}$
- (c)  $v_2^2 - v_1^2 = 2gh$




(d) The energy per unit mass of the liquid is the same in both sections of the tube

38. A sniper fires a rifle bullet into a gasoline tank making a hole 53.0 m below the surface of gasoline. The tank was sealed at 3.10 atm. The stored gasoline has a density of  $660 \text{ kgm}^{-3}$ . The velocity with which gasoline begins to shoot out of the hole is

- (a)  $27.8 \text{ ms}^{-1}$                               (b)  $41.0 \text{ ms}^{-1}$
- (c)  $9.6 \text{ ms}^{-1}$                               (d)  $19.7 \text{ ms}^{-1}$

39. An L-shaped tube with a small orifice is held in a water stream as shown in fig. The upper end of the tube is 10.6 cm above the surface of water. What will be the height of the jet of water coming from the orifice? Velocity of water stream is  $2.45 \text{ m/s}$


- (a) Zero
- (b) 20.0 cm
- (c) 10.6 cm
- (d) 40.0 cm



40. Fig. represents vertical sections of four wings moving horizontally in air. In which case the force is upwards

- (a)
- (b)
- (c)
- (d)

41. An L-shaped glass tube is just immersed in flowing water such that its opening is pointing against flowing water.




If the speed of water current is  $v$ , then

- (a) The water in the tube rises to height  $\frac{v^2}{2g}$   
 (b) The water in the tube rises to height  $\frac{g}{2v^2}$   
 (c) The water in the tube does not rise at all  
 (d) None of these


42. A tank is filled with water up to a height  $H$ . Water is allowed to come out of a hole  $P$  in one of the walls at a depth  $D$  below the surface of water. Express the horizontal distance  $x$  in terms of  $H$  and  $D$  [MNR 1992; CPMT 2004]

- (a)  $x = \sqrt{D(H-D)}$   
 (b)  $x = \sqrt{\frac{D(H-D)}{2}}$   
 (c)  $x = 2\sqrt{D(H-D)}$   
 (d)  $x = 4\sqrt{D(H-D)}$



43. A cylindrical vessel of 90 cm height is kept filled upto the brim. It has four holes 1, 2, 3, 4 which are respectively at heights of 20 cm, 30 cm, 45 cm and 50 cm from the horizontal floor  $PQ$ . The water falling at the maximum horizontal distance from the vessel comes from [CPMT 1989]

- (a) Hole number 4  
 (b) Hole number 3  
 (c) Hole number 2  
 (d) Hole number 1



44. A rectangular vessel when full of water takes 10 minutes to be emptied through an orifice in its bottom. How much time will it take to be emptied when half filled with water

- (a) 9 minute (b) 7 minute  
 (c) 5 minute (d) 3 minute

45. A streamlined body falls through air from a height  $h$  on the surface of a liquid. If  $d$  and  $D$  ( $D > d$ ) represents the densities of the material of the body and liquid respectively, then the time after which the body will be instantaneously at rest, is

- (a)  $\sqrt{\frac{2h}{g}}$  (b)  $\sqrt{\frac{2h \cdot D}{g \cdot d}}$

- (c)  $\sqrt{\frac{2h \cdot d}{g \cdot D}}$  (d)  $\sqrt{\frac{2h}{g} \left( \frac{d}{D-d} \right)}$

46. A large tank is filled with water to a height  $H$ . A small hole is made at the base of the tank. It takes  $\tau_1$  time to decrease the height of water to  $\frac{H}{\eta}$  ( $\eta > 1$ ); and it takes  $\tau_2$  time to take out the rest

of water. If  $\tau_1 = \tau_2$ , then the value of  $\eta$  is

- (a) 2 (b) 3  
 (c) 4 (d)  $2\sqrt{2}$

47. Velocity of water in a river is [CBSE PMT 1988]

- (a) Same everywhere  
 (b) More in the middle and less near its banks  
 (c) Less in the middle and more near its banks  
 (d) Increase from one bank to other bank

48. As the temperature of water increases, its viscosity

- (a) Remains unchanged  
 (b) Decreases  
 (c) Increases  
 (d) Increases or decreases depending on the external pressure

49. The coefficient of viscosity for hot air is

- (a) Greater than the coefficient of viscosity for cold air  
 (b) Smaller than the coefficient of viscosity for cold air  
 (c) Same as the coefficient of viscosity for cold air  
 (d) Increases or decreases depending on the external pressure

50. A good lubricant should have

- (a) High viscosity (b) Low viscosity  
 (c) Moderate viscosity (d) High density

51. We have three beakers  $A$ ,  $B$  and  $C$  containing glycerine, water and kerosene respectively. They are stirred vigorously and placed on a table. The liquid which comes to rest at the earliest is

- (a) Glycerine  
 (b) Water  
 (c) Kerosene

- (d) All of them at the same time
52. A small drop of water falls from rest through a large height  $h$  in air; the final velocity is
- (a)  $\propto \sqrt{h}$   
 (b)  $\propto h$   
 (c)  $\propto (1/h)$   
 (d) Almost independent of  $h$
53. The rate of flow of liquid in a tube of radius  $r$ , length  $l$ , whose ends are maintained at a pressure difference  $P$  is  $V = \frac{\pi Q P r^4}{\eta l}$  where  $\eta$  is coefficient of the viscosity and  $Q$  is
- [DCE 2002]
- (a) 8  
 (b)  $\frac{1}{8}$   
 (c) 16  
 (d)  $\frac{1}{16}$
54. In Poiseuille's method of determination of coefficient of viscosity, the physical quantity that requires greater accuracy in measurement is
- [EAMCET 2001]
- (a) Pressure difference  
 (b) Volume of the liquid collected  
 (c) Length of the capillary tube  
 (d) Inner radius of the capillary tube
55. Two capillary tubes of the same length but different radii  $r_1$  and  $r_2$  are fitted in parallel to the bottom of a vessel. The pressure head is  $P$ . What should be the radius of a single tube that can replace the two tubes so that the rate of flow is same as before
- (a)  $r_1 + r_2$   
 (b)  $r_1^2 + r_2^2$   
 (c)  $r_1^4 + r_2^4$   
 (d) None of these
56. Two capillaries of same length and radii in the ratio 1 : 2 are connected in series. A liquid flows through them in streamlined condition. If the pressure across the two extreme ends of the combination is 1 m of water, the pressure difference across first capillary is
- (a) 9.4 m  
 (b) 4.9 m  
 (c) 0.49 m  
 (d) 0.94 m
57. Water flows in a streamlined manner through a capillary tube of radius  $a$ , the pressure difference being  $P$  and the rate of flow  $Q$ . If the radius is reduced to  $a/2$  and the pressure increased to  $2P$ , the rate of flow becomes
- (a)  $4Q$   
 (b)  $Q$   
 (c)  $\frac{Q}{4}$   
 (d)  $\frac{Q}{8}$
58. A viscous fluid is flowing through a cylindrical tube. The velocity distribution of the fluid is best represented by the diagram
- [BCECE 2005]
- 
- (a) (b)  
 (c) (d) None of these
59. Water is flowing in a pipe of diameter 4 cm with a velocity 3 m/s. The water then enters into a tube of diameter 2 cm. The velocity of water in the other pipe is
- [BCECE 2005]
- (a) 3 m/s  
 (b) 6 m/s  
 (c) 12 m/s  
 (d) 8 m/s
60. Two capillary of length  $L$  and  $2L$  and of radius  $R$  and  $2R$  are connected in series. The net rate of flow of fluid through them will be (given rate of the flow through single capillary,  $X = \pi P R^4 / 8 \eta L$ )
- [DCE 2005]
- (a)  $\frac{8}{9} X$   
 (b)  $\frac{9}{8} X$   
 (c)  $\frac{5}{7} X$   
 (d)  $\frac{7}{5} X$
61. When a body falls in air, the resistance of air depends to a great extent on the shape of the body, 3 different shapes are given. Identify the combination of air resistances which truly represents the physical situation. (The cross sectional areas are the same).
- [KCEET 2005]
- 
- (1) Disc (2) Ball (3) Cigar shaped

- (a)  $1 < 2 < 3$                       (b)  $2 < 3 < 1$   
 (c)  $3 < 2 < 1$                       (d)  $3 < 1 < 2$

62. Water falls from a tap, down the streamline  
 [Orissa JEE 2005]

- (a) Area decreases              (b) Area increases  
 (c) Velocity remains same (d) Area remains same

63. A manometer connected to a closed tap reads  $4.5 \times 10^5$  pascal. When the tap is opened the reading of the manometer falls to  $4 \times 10^5$  pascal. Then the velocity of flow of water is  
 [Kerala PET 2005]

- (a)  $7 \text{ ms}^{-1}$                       (b)  $8 \text{ ms}^{-1}$   
 (c)  $9 \text{ ms}^{-1}$                       (d)  $10 \text{ ms}^{-1}$

64. What is the velocity  $v$  of a metallic ball of radius  $r$  falling in a tank of liquid at the instant when its acceleration is one-half that of a freely falling body ? (The densities of metal and of liquid are  $\rho$  and  $\sigma$  respectively, and the viscosity of the liquid is  $\eta$ ).  
 [Kerala PET 2005]

- (a)  $\frac{r^2 g}{9\eta}(\rho - 2\sigma)$               (b)  $\frac{r^2 g}{9\eta}(2\rho - \sigma)$   
 (c)  $\frac{r^2 g}{9\eta}(\rho - \sigma)$               (d)  $\frac{2r^2 g}{9\eta}(\rho - \sigma)$

65. Consider the following equation of Bernoulli's theorem.  $P + \frac{1}{2}\rho v^2 + \rho gh = K$  (constant)


The dimensions of  $K/P$  are same as that of which of the following [AFMC 2005]

- (a) Thrust                      (b) Pressure  
 (c) Angle                      (d) Viscosity

66. An incompressible fluid flows steadily through a cylindrical pipe which has radius  $2r$  at point  $A$  and radius  $r$  at  $B$  further along the flow direction. If the velocity at point  $A$  is  $v$ , its velocity at point  $B$  is  
 [Kerala PMT 2005]


- (a)  $2v$                       (b)  $v$   
 (c)  $v/2$                       (d)  $4v$

1. A U-tube in which the cross-sectional area of the limb on the left is one quarter, the limb on the right contains mercury (density  $13.6 \text{ g/cm}^3$ ). The level of mercury in the narrow limb is at a distance of  $36 \text{ cm}$  from the upper end of the tube. What will be the rise in the level of mercury in the right limb if the left limb is filled to the top with water



- (a)  $1.2 \text{ cm}$   
 (b)  $2.35 \text{ cm}$   
 (c)  $0.56 \text{ cm}$   
 (d)  $0.8 \text{ cm}$

2. A homogeneous solid cylinder of length  $L$  ( $L < H/2$ ). Cross-sectional area  $A/5$  is immersed such that it floats with its axis vertical at the liquid-liquid interface with length  $L/4$  in the denser liquid as shown in the fig. The lower density liquid is open to atmosphere having pressure  $P_0$ . Then density  $D$  of solid is given by  
 [IIT-JEE 1995]




- (a)  $\frac{5}{4}d$   
 (b)  $\frac{4}{5}d$   
 (c)  $d$   
 (d)  $\frac{d}{5}$

3. A wooden block, with a coin placed on its top, floats in water as shown in fig. the distance  $l$  and  $h$  are shown there. After some time the coin falls into the water. Then

[IIT-JEE (Screening) 2002]

- (a)  $l$  decreases and  $h$  increases  
 (b)  $l$  increases and  $h$  decreases  
 (c) Both  $l$  and  $h$  increase  
 (d) Both  $l$  and  $h$  decrease




4. A vessel contains oil (density =  $0.8 \text{ gm/cm}^3$ ) over mercury (density =  $13.6 \text{ gm/cm}^3$ ). A homogeneous sphere floats with half of its volume immersed in mercury and the other half

in oil. The density of the material of the sphere in  $gm/cm^3$  is [IIT-JEE 1988]

- (a) 3.3 (b) 6.4  
(c) 7.2 (d) 12.8

5. A body floats in a liquid contained in a beaker. The whole system as shown falls freely under gravity. The upthrust on the body due to the liquid is



[IIT-JEE 1982]


- (a) Zero  
(b) Equal to the weight of the liquid displaced  
(c) Equal to the weight of the body in air  
(d) Equal to the weight of the immersed position of the body

6. A liquid is kept in a cylindrical vessel which is being rotated about a vertical axis through the centre of the circular base. If the radius of the vessel is  $r$  and angular velocity of rotation is  $\omega$ , then the difference in the heights of the liquid at the centre of the vessel and the edge is

- (a)  $\frac{r\omega}{2g}$  (b)  $\frac{r^2\omega^2}{2g}$   
(c)  $\sqrt{2gr\omega}$  (d)  $\frac{\omega^2}{2gr^2}$

7. Water is filled in a cylindrical container to a height of  $3m$ . The ratio of the cross-sectional area of the orifice and the beaker is  $0.1$ . The square of the speed of the liquid coming out from the orifice is ( $g = 10 m/s^2$ ) [IIT JEE 2004]

- (a)  $50 m^2/s^2$   
(b)  $50.5 m^2/s^2$   
(c)  $51 m^2/s^2$   
(d)  $52 m^2/s^2$




8. A large open tank has two holes in the wall. One is a square hole of side  $L$  at a depth  $y$  from the top and the other is a circular hole of radius  $R$  at a depth  $4y$  from the top. When the tank is

completely filled with water the quantities of water flowing out per second from both the holes are the same. Then  $R$  is equal to [IIT-JEE (Screening) 2000]

- (a)  $2\pi L$  (b)  $\frac{L}{\sqrt{2\pi}}$   
(c)  $L$  (d)  $\frac{L}{2\pi}$


9. A cylinder containing water up to a height of  $25 cm$  has a hole of cross-section  $\frac{1}{4} cm^2$  in its bottom. It is counterpoised in a balance. What is the initial change in the balancing weight when water begins to flow out

- (a) Increase of  $12.5 gm-wt$   
(b) Increase of  $6.25 gm-wt$   
(c) Decrease of  $12.5 gm-wt$   
(d) Decrease of  $6.25 gm-wt$




10. There are two identical small holes of area of cross-section  $a$  on the opposite sides of a tank containing a liquid of density  $\rho$ . The difference in height between the holes is  $h$ . Tank is resting on a smooth horizontal surface. Horizontal force which will have to be applied on the tank to keep it in equilibrium is

- (a)  $gh\rho a$   
(b)  $\frac{2gh}{\rho a}$   
(c)  $2\rho agh$   
(d)  $\frac{\rho gh}{a}$



11. Two communicating vessels contain mercury. The diameter of one vessel is  $n$  times larger than the diameter of the other. A column of water of height  $h$  is poured into the left vessel. The mercury level will rise in the right-hand



- (a)  $\frac{r^2 h}{(n+1)^2 s}$                       (b)  $\frac{h}{(r^2 + 1) s}$   
 (c)  $\frac{h}{(n+1)^2 s}$                       (d)  $\frac{h}{r^2 s}$

12. A uniform rod of density  $\rho$  is placed in a wide tank containing a liquid of density  $\rho_0 (\rho_0 > \rho)$ . The depth of liquid in the tank is half the length of the rod. The rod is in equilibrium, with its lower end resting on the bottom of the tank. In this position the rod makes an angle  $\theta$  with the horizontal

- (a)  $\sin \theta = \frac{1}{2} \sqrt{\rho_0 / \rho}$                       (b)  $\sin \theta = \frac{1}{2} \cdot \frac{\rho_0}{\rho}$   
 (c)  $\sin \theta = \sqrt{\rho / \rho_0}$                       (d)  $\sin \theta = \rho_0 / \rho$

13. A block of ice floats on a liquid of density 1.2 in a beaker then level of liquid when ice completely melt

[IIT-JEE 1994]

- (a) Remains same                      (b) Rises  
 (c) Lowers                      (d) (a), (b) or (c)



14. A vessel of area of cross-section  $A$  has liquid to a height  $H$ . There is a hole at the bottom of vessel having area of cross-section  $a$ . The time taken to decrease the level from  $H_1$  to  $H_2$  will be



- (a)  $\frac{A}{a} \sqrt{\frac{2}{g}} [\sqrt{H_1} - \sqrt{H_2}]$                       (b)  $\sqrt{2gh}$   
 (c)  $\sqrt{2gh(H_1 - H_2)}$                       (d)  $\frac{A}{a} \sqrt{\frac{g}{2}} [\sqrt{H_1} - \sqrt{H_2}]$

## Graphical Questions


1. A lead shot of 1mm diameter falls through a long column of glycerine. The variation of its velocity  $v$ , with distance covered is represented by

[AIIMS 2003]

- (a)                       (b) 


- (c)                       (d) 





2. A small spherical solid ball is dropped from a great height in a viscous liquid. Its journey in the liquid is best described in the diagram given below by the




- (a) Curve A  
 (b) Curve B  
 (c) Curve C  
 (d) Curve D



3. The diagram shows a cup of tea seen from above. The tea has been stirred and is now rotating without turbulence. A graph showing the speed  $v$  with which the liquid is crossing points at a distance  $x$  from  $O$  along a radius  $XO$  would look




- (a)                       (b)   
 (c)                       (d) 


4. Water flows through a frictionless duct with a cross-section varying as shown in fig. Pressure  $p$  at points along



- (a)                       (b) 



5. From amongst the following curves, which one shows the variation of the velocity  $v$  with time  $t$  for a small sized spherical body falling vertically in a long column of a viscous liquid  
[CPMT 1990]



## Assertion & Reason

For AIIMS Aspirants

Read the assertion and reason carefully to mark the correct option out of the options given below:

- (a) If both assertion and reason are true and the reason is the correct explanation of the assertion.  
 (b) If both assertion and reason are true but reason is not the correct explanation of the assertion.  
 (c) If assertion is true but reason is false.  
 (d) If the assertion and reason both are false.  
 (e) If assertion is false but reason is true.

1. Assertion : Pascal law is the working principle of a hydraulic lift.  
 Reason : Pressure is equal to thrust per area.  
 2. Assertion : The blood pressure in humans is greater at the feet than at the brain.  
 Reason : Pressure of liquid at any point is proportional to height, density of

liquid and acceleration due to gravity.

3. Assertion : Hydrostatic pressure is a vector quantity.  
 Reason : Pressure is force divided by area, and force is a vector quantity.  
 4. Assertion : To float, a body must displace liquid whose weight is greater than the actual weight of the body.  
 Reason : The body will experiences no net downward force, in the case of floating.  
 5. Assertion : A man sitting in a boat which is floating on a pond. If the man drinks some water from the pond, the level of the water in the pond decreases.  
 Reason : According to Archimede's principle the weight displaced by body is equal to the weight of the body.  
 6. Assertion : A piece of ice floats in water, the level of water remains unchanged when the ice melts completely.  
 Reason : According to Archimede's principle, the loss in weight of the body in the liquid is equal to the weight of the liquid displaced by the immersed part of the body.  
 7. Assertion : The velocity increases, when water flowing in broader pipe enter a narrow pipe.  
 Reason : According to equation of continuity, product of area and velocity is constant.  
 8. Assertion : The velocity of fall of a man jumping with a parachute first increases and then becomes constant.  
 Reason : The constant velocity of fall of man is called terminal velocity.  
 9. Assertion : The velocity of flow of a liquid is smaller when pressure is larger and viceversa.



- Reason : According to Bernoulli's theorem, for the stream line flow of an ideal liquid, the total energy per unit mass remains constant.
10. Assertion : The shape of an automobile is so designed that its front resembles the stream line pattern of the fluid through which it moves.  
Reason : The resistance offered by the fluid is maximum.
11. Assertion : The size of the needle of a syringe controls flow rate better than the thumb pressure exerted by a doctor while administering an injection.  
Reason : Flow rate is independent of pressure exerted by the thumb of the doctor.
12. Assertion : A fluid flowing out of a small hole in a vessel apply a backward thrust on the vessel.  
Reason : According to equation of continuity, the product of area and velocity remain constant.
13. Assertion : For a floating body to be in stable equilibrium, its centre of buoyancy must be located above the centre of gravity.  
Reason : The torque produced by the weight of the body and the upthrust will restore body back to its normal position, after the body is disturbed.
14. Assertion : Water flows faster than honey.  
Reason : The coefficient of viscosity of water is less than honey.
15. Assertion : The viscosity of liquid increases rapidly with rise of temperature.  
Reason : Viscosity of a liquid is the property of the liquid by virtue of which it opposes the relative motion amongst its different layers.
16. Assertion : Aeroplanes are made to run on the runway before take off, so that they acquire the necessary lift.  
Reason : According to Bernoulli's theorem, as velocity increases pressure decreases and viceversa.
17. Assertion : Sudden fall of pressure at a place indicates storm.  
Reason : Air flows from higher pressure to lower pressure.
18. Assertion : Machine parts are jammed in winter.  
Reason : The viscosity of lubricant used in machine parts increase at low temperature.
19. Assertion : A block of wood is floating in a tank containing water. The apparent weight of the floating block is equal to zero.  
Reason : Because the entire weight of the block is supported by the buoyant force (the upward thrust) due to water.
20. Assertion : A rain drop after falling through some height attains a constant velocity.  
Reason : At constant velocity, the viscous drag is just equal to its weight.
21. Assertion : paper pins are made to have pointed end.  
Reason : Because pointed pins have very small area due to which even for small applied force it exert large pressure on the surface.
22. Assertion : Railways tracks are laid on small sized wooden sleepers.  
Reason : Small sized wooden sleepers are used so that rails exert more pressure on the railway track. Due to which rail does not leave the track
23. Assertion : It is difficult to stop bleeding from a cut in the body at high altitudes.  
Reason : The atmospheric pressure at high altitude is lesser than the blood pressure.

24. Assertion : To empty an oil tank, two holes are made.  
Reason : Oil will come out two holes so it will emptied faster.
25. Assertion : Terminal velocity is same as the critical velocity.  
Reason : The constant velocity of fall of a body through a viscous fluid is called terminal velocity.
26. Assertion : When two boats sails parallel in the same direction and close to each other, they are pulled towards each other.  
Reason : The viscous drag on a spherical body moving with speed  $v$  is proportional to  $v$ .
27. Assertion : Cars and aeroplanes are streamlined.  
Reason : This is done to reduce the backward drag due to atmosphere.
28. Assertion : Bernoulli's theorem holds for incompressible, non-viscous fluids.  
Reason : The factor  $\frac{v^2}{2g}$  is called velocity head.

# Answers

## Pressure and Density

1	b	2	c	3	b	4	b	5	c
6	b	7	c	8	b	9	d	10	c
11	c	12	c	13	a	14	d	15	a
16	b	17	a	18	b	19	d	20	b
21	c	22	a	23	b	24	b	25	b
26	b	27	d	28	a	29	c	30	c
31	d								

## Pascal's Law and Archimides Principle

1	c	2	a	3	c	4	c	5	b
6	d	7	a	8	b	9	bc	10	c

11	c	12	d	13	a	14	b	15	b
16	d	17	a	18	a	19	b	20	b
21	c	22	c	23	b	24	b	25	b
26	d	27	a						

## Fluid Flow

1	b	2	a	3	c	4	d	5	d
6	a	7	d	8	b	9	a	10	c
11	b	12	a	13	c	14	a	15	d
16	b	17	b	18	c	19	c	20	b
21	d	22	b	23	a	24	d	25	c
26	a	27	a	28	b	29	a	30	a
31	a	32	b	33	c	34	a	35	b
36	ad	37	acd	38	b	39	b	40	a
41	a	42	c	43	b	44	b	45	d
46	c	47	b	48	b	49	a	50	a
51	a	52	d	53	b	54	d	55	d
56	d	57	d	58	c	59	c	60	a
61	c	62	a	63	d	64	c	65	c
66	d								

## Critical Thinking Questions

1	c	2	a	3	d	4	c	5	a
6	b	7	a	8	b	9	c	10	c
11	b	12	a	13	b	14	a		

## Graphical Questions

1	a	2	b	3	d	4	a	5	d
---	---	---	---	---	---	---	---	---	---

## Assertion and Reason

1	b	2	a	3	e	4	c	5	e
6	a	7	a	8	b	9	a	10	c
11	c	12	a	13	a	14	a	15	e
16	a	17	a	18	a	19	a	20	a
21	a	22	d	23	a	24	c	25	e
26	b	27	a	28	b				